



## Plague FAQs

### **What is plague?**

Plague is a severe disease caused by the bacteria *Yersinia pestis* (*Y. pestis*). Plague is usually carried by rodents but can be spread to man by fleas. The disease is rare in the United States, but occasional cases do occur in the western region of the country. Plague has occasionally occurred in the Panhandle of Oklahoma.

### **Why are we concerned about pneumonic plague as a bioweapon?**

*Yersinia pestis* used in an aerosol attack could cause cases of pneumonic form of plague. One to six days after becoming infected with bacteria, people would develop pneumonic plague. Once people have the disease, the bacteria can spread to others who have close contact with them. Because of the delay between being exposed to the bacteria and becoming sick, people could travel over large area before becoming contagious and possibly infecting others. Controlling the disease would then be more difficult. A bioweapon carrying *Y. pestis* is possible because the bacterium occurs in nature and could be isolated and grown in quantity in a laboratory. Even so, manufacturing an effective weapon using *Y. pestis* would require advanced knowledge and technology.

### **Is pneumonic plague different from bubonic plague?**

Yes. Both are caused by *Yersinia pestis*, but they are transmitted differently and their symptoms differ. Pneumonic plague can be transmitted from person to person; bubonic plague cannot. Pneumonic plague affects the lungs and is transmitted when a person breathes in *Y. pestis* particles in the air. Bubonic plague is transmitted through the bite of an infected flea or exposure to infected material through a break in the skin. If bubonic plague is not treated, however, the bacteria can spread through the bloodstream and infect the lungs, causing a secondary case of pneumonic plague.

*For further information, contact the Oklahoma City-County Health Department  
(405) 425-4437*



## Plague FAQs

### **How do people become infected with pneumonic plague?**

Pneumonic plague occurs when *Yersinia pestis* infects the lungs. Transmission can take place if someone breathes in *Y. pestis* particles, which could happen in an aerosol release during a bioterrorism attack. Pneumonic plague is also transmitted by breathing in *Y. pestis* suspended in respiratory droplets from a person (or animal) with pneumonic plague. Respiratory droplets are spread most readily by coughing or sneezing. Becoming infected in this way usually requires direct and close (within 6 feet) contact with the ill person or animal.

### **What are the signs and symptoms of pneumonic plague?**

Patients usually have fever, weakness, and rapidly developing pneumonia with shortness of breath, chest pain, cough, and sometimes bloody or watery sputum. Nausea, vomiting, and abdominal pain may also occur. Without early treatment, pneumonic plague usually leads to respiratory failure, shock, and rapid death.

### **Does plague occur naturally?**

Yes. The World Health Organization reports 1,000 to 3,000 cases of plague worldwide every year. An average of 5 to 15 cases occur each year in the western United States. These cases are scattered and occur in rural or semi-rural areas. Most cases are of the bubonic form of the disease. Naturally occurring pneumonic plague is uncommon, although small outbreaks do occur. Both types of plague are readily controlled by standard public health response measures.

### **Can a person exposed to pneumonic plague avoid becoming sick?**

Yes. People who have had close contact with an infected person can greatly reduce the chance of becoming sick if they begin treatment within 7 days of their exposure. Treatment consists of taking antibiotics for at least 7 days.

*For further information, contact the Oklahoma City-County Health Department  
(405) 425-4437*

revised 05/2013

**WELLNESS NOW**



## Plague FAQs

### **How quickly would someone get sick if exposed to plague bacteria through the air?**

Someone exposed to *Yersinia pestis* through the air – either from intentional aerosol release or from close and direct exposure to someone with plague pneumonia – would become ill within 1 to 6 days.

### **Can pneumonic plague be treated?**

Yes. To prevent a high risk of death, antibiotics should be given within 24 hours of the first symptoms. Several types of antibiotics are effective for curing the disease and for preventing it. Available oral medications are a tetracycline (such as doxycycline) or a fluoropuinolone (such as ciprofloxacin). For injection or intravenous use, streptomycin or gentamycin antibiotics are used. Early in the response to a bioterrorism attack, these drugs would be tested to determine which is most effective against the particular weapon that was used.

### **Would enough medication be available in the event of a bioterrorism attack involving pneumonic plague?**

National and state public health officials have large supplies of drugs needed in the event of a bioterrorism attack. These supplies can be sent anywhere in the United States within 12 hours.

*For further information, contact the Oklahoma City-County Health Department  
(405) 425-4437*

revised 05/2013

**WELLNESS NOW**



## Plague FAQs

### **What should someone do if they suspect they or others have been exposed to plague?**

Get immediate medical attention: To prevent illness, a person who has been exposed to pneumonic plague must receive antibiotic treatment without delay. If an exposed person becomes ill, antibiotics must be administered within 24 hours of their first symptoms to reduce the risk of death. Notify authorities: Immediately notify local and state health departments so they can begin to investigate and control the problem right away. If bioterrorism is suspected, the health departments will notify the CDC, FBI, and other appropriate authorities.

### **How can the general public reduce the risk of getting pneumonic plague from another person or giving it to someone else?**

If possible, avoid close contact with other people. People having direct and close contact with someone with pneumonic plague should wear tightly fitting disposable surgical masks. If surgical masks are not available, even makeshift face coverings made of layers of cloth may be helpful in an emergency. People who have been exposed to a contagious person can be protected from developing plague by receiving prompt antibiotic treatment.

### **How is plague diagnosed?**

The first step is evaluation by a health worker. If the health worker suspects pneumonic plague, samples of the patient's blood, sputum, or lymph node aspirate are sent to a laboratory for testing. Once the laboratory receives the sample, preliminary results can be ready in less than two hours. Confirmation will take longer, usually 24 to 48 hours.

*For further information, contact the Oklahoma City-County Health Department  
(405) 425-4437*

revised 05/2013

**WELLNESS NOW**



## Plague FAQs

### **How long can plague bacteria exist in the environment?**

*Yersinia pestis* is easily destroyed by sunlight and drying. Even so, when released into air, the bacterium will survive for up to one hour, depending on conditions.

### **Is a vaccine available to prevent pneumonic plague?**

Currently, no plague vaccine is available in the United States. Research is in progress, but we are not likely to have vaccines for several years or more.

*For further information, contact the Oklahoma City-County Health Department  
(405) 425-4437*

revised 05/2013

**WELLNESS NOW**