



Measles Fact Sheet

What is measles?

Measles, not to be confused with German measles (rubella), is an extremely contagious disease caused by a virus. A high fever and a red rash covering most of the body develops in almost every case. Measles can also cause pneumonia, deafness, and inflammation of the brain and death.

Who gets measles?

Anyone, regardless of age, who has not had measles or has not been adequately immunized may get the disease. Most cases occur in unimmunized preschoolers and young adults.

How is measles spread?

Measles is spread through the air by droplets from the nose, throat, and mouth of an infected person by coughing, sneezing, or simply talking.

What are the symptoms of measles?

Measles causes a high fever, cough, runny nose, watery eyes, and a red rash moving from the face to the rest of the body. The symptoms may last from 1 to 2 weeks.

How long is an infected person able to spread measles?

An infected person is able to spread measles from 4 days before the rash starts to 4 days after the rash appears.

OCCHD.ORG |  [Facebook](#) - Search OKC-County Health or @OCCHD

[OCCHD App](#) - Search OCCHD on the Apple Store



or [Google Play](#)





Measles Fact Sheet

Can a person get measles again?

No. One attack of measles provides protection for life.

What is the treatment for measles?

Bedrest is advisable while the patient has a fever. Infants with very high fever and children with headaches should be treated with anti-fever medicines.

Should people who have been around a person infected with measles be treated?

Anyone who has not had measles or has not been immunized should receive measles vaccine immediately. Infants under one year old may receive immune globulin (IG) or measles vaccine depending on the child's age.

How can the spread of measles be stopped?

The best way to stop measles is to immunize all children by 15 months of age with measles vaccine (combined with rubella and mumps vaccines or MMR). A second dose of measles vaccine between 4 and 6 years of age or before starting kindergarten is now required for children entering school in Oklahoma. Adults born after 1956 may also need measles vaccine.

Does measles vaccine cause reactions?

About 1 child in 5 will develop a rash or fever which begins a week or two after vaccination and lasts a few days. On rare occasions (1 in every million shots) a

OCCHD.ORG |  [Facebook](#) - Search OKC-County Health or @OCCHD

[OCCHD App](#) - Search OCCHD on the Apple Store



or [Google Play](#)





Measles Fact Sheet

child may have a more serious reaction to MMR vaccine such as inflammation of the brain (encephalitis).

Where can children receive measles vaccine?

Children and adults may receive measles vaccine at county health departments throughout Oklahoma or check with your private doctor.

*For further information, contact the
Oklahoma City-County Health
Department (405) 425-4437*

revised 04/2013

OCCHD.ORG |  Facebook - Search OKC-County Health or @OCCHD

OCCHD App - Search OCCHD on the Apple Store  or Google Play 



OCCHD.ORG |  **Facebook** - Search OKC-County Health or @OCCHD

OCCHD App - Search OCCHD on the Apple Store  or Google Play 